

Quenched Kosterlitz-Thouless Superfluid Transitions

Han-Ching Chu and Gary A. Williams

Department of Physics and Astronomy, University of California, Los Angeles, CA 90095

(Dated: February 1, 2008)

Rapidly quenched Kosterlitz-Thouless (KT) superfluid transitions are studied by solving the Fokker-Planck equation for the vortex-pair dynamics in conjunction with the KT recursion relations. Power-law decays of the vortex density at long times are found, and the results are in agreement with a scaling proposal made by Minnhagen and co-workers for the dynamical critical exponent. The superfluid density is strongly depressed after a quench, with the subsequent recovery being logarithmically slow for starting temperatures near T_{KT} . No evidence is found of vortices being "created" in a rapid quench, there is only decay of the existing thermal vortex pairs.

PACS numbers: 64.60.Ht, 67.40.Fd, 67.40.Vs, 05.70.Ln

There is considerable current interest in the properties of quenched superfluid transitions, since they may have relevance to cosmic-string phase transitions in the rapidly-cooling early universe [1]. Experiments [2] have been carried out in superfluid ^4He attempting to observe vortex structures "created" in rapid pressure quenches through the transition, but no vortices have been observed, which is a puzzle [3].

To try to understand these quenched transitions we have constructed [4] an analytic formulation of the quenched Kosterlitz-Thouless (KT) superfluid transition in two dimensions (2D). Although not the three-dimensional (3D) realm of the experiments, the simplicity of the KT vortex renormalization scheme allows insights into the vortex dynamics that should be useful in extending the calculations to 3D [5]. Previous studies of quenched 2D transitions have included computer simulations [6, 7, 8] and dynamic scaling theories [9, 10, 11], but there is still considerable controversy over the physics involved. The various dynamic scaling theories give quite different predictions for the temperature dependence of the dynamic scaling exponent z , and Bray and co-workers [10] have even suggested that there may be violations of dynamic scaling. Unfortunately the computer simulations are not able to resolve the questions in the field, since it has recently been shown [8] that different boundary conditions in the simulations give different results for the dynamics.

We consider a thin superfluid helium film on a flat substrate, which is coupled to a thermal reservoir at temperature T . The vortex pairs in the film are characterized by the distribution function $\Gamma(r, t)$, which is the density of pairs of separation r and $r+dr$. It is determined from the 2D Fokker-Planck equation [12],

$$\frac{\partial \Gamma}{\partial t} = \frac{2D}{a_o^2} \frac{\partial}{\partial r'} \cdot \left(\frac{\partial \Gamma}{\partial r'} + \Gamma \frac{\partial}{\partial r'} \left(\frac{U}{k_B T} \right) \right) , \quad (1)$$

where $r' = r/a_o$ with a_o the vortex core radius, U is the pair interaction energy, and D is the diffusion coefficient characterizing the mutual friction drag force on the vortex cores of a pair. It can be seen that the time in this equation is

scaled by $\tau_o = a_o^2/2D$, the diffusion time of the smallest pairs of separation a_o . In common with the scaling theories and simulations, we assume that any renormalization of D can be neglected. At equilibrium where $\frac{\partial \Gamma}{\partial t} = 0$ the solution of Eq. (1) is just the usual Boltzmann distribution, $\Gamma = a_o^{-4} \exp(-(U + 2E_c)/k_B T)$ where $2E_c$ is the core energy of a pair.

The vortex interaction energy U in Eq. (1) is determined using the Kosterlitz-Thouless (KT) vortex renormalization methods [13]. The Kosterlitz recursion relations can be written in the form

$$\frac{1}{k_B T} \frac{\partial U}{\partial r} = \frac{2\pi K}{r} \quad (2)$$

and

$$\frac{\partial K}{\partial r} = -4\pi^3 r^3 K^2 \Gamma , \quad (3)$$

where $K = \hbar^2 \sigma_s / m^2 k_B T$ is the dimensionless areal superfluid density. We take the core energy to be the Villain-model value $2E_c/k_B T = \pi^2 K_o$, where K_o is the starting value of K at $r = a_o$. In thermal equilibrium Eqs. (1–3) lead to the well-known result that the superfluid density has a universal jump to zero just above the transition temperature T_{KT} [14].

To study the quenched transition [4] the film is first equilibrated with the heat bath at temperature T , generating an equilibrium distribution of vortex pairs. The top curve in Fig. 1 at $t = 0$ is the distribution at T_{KT} ; it varies asymptotically as $(r/a_o)^{-2\pi K}$. The temperature of the heat bath is then reduced abruptly to a low temperature, $0.1 T_{KT}$, where in equilibrium the vortex density is over 20 orders of magnitude smaller. However, due to the Kelvin circulation theorem the pairs cannot suddenly disappear, since the only way they can be extracted by the heat bath is if the plus-minus pairs annihilate at $r = a_o$, where the remaining core energy is converted to phonons. This can take quite a long time to occur, however, since the pairs only slowly lose kinetic energy to the diffusive frictional force of the heat bath as they move together towards annihilation [15].

To solve for the time dependence of the distribution function after the quench we combine Eqs. (1) and (2), which become

$$\frac{\partial \Gamma}{\partial t} = \frac{2D}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \Gamma}{\partial r} + 2\pi K \Gamma \right) \quad . \quad (4)$$

We solve this in conjunction with Eq. (3) using the method of lines with a third-order Runge-Kutta technique on a finite domain, to a maximum pair separation $R/a_o = e^{10} = 2.2 \times 10^4$. The boundary conditions used are that the flux of pairs $J = -2D(r \frac{\partial \Gamma}{\partial r} + 2\pi K \Gamma)$ be continuous across the boundary $r = R$, and that Γ drop abruptly to zero at $r = a_o$, which is equivalent to putting a delta-function sink term at a_o on the right-hand side of Eq. (4) to generate the annihilation at that point.




FIG. 1: Time dependence (in units of the diffusion time τ_o) of the pair distribution function Γ (in units a_o^{-4}) for an instantaneous quench from T_{KT} to $0.1 T_{KT}$.

Figure 1 shows the time dependence of the distribution function for an instantaneous quench to $0.1 T_{KT}$ from a starting temperature barely below T_{KT} (from $K_o = 0.747853$, compared to the critical value 0.747852 at T_{KT}). As expected for the annihilation process the smallest pairs decay away first, in the classic pattern of a phase-ordering transition where at long times the largest scales become more dominant. By integrating the distribution function over d^2r the pair density is obtained, shown as the top curve in Fig. 2a. At long times the density decreases nearly as t^{-1} , in agreement with the scaling theories [9, 10, 11] at T_{KT} . We find, however, that the slope only approaches -1 logarithmically slowly, being -1.12 at $t = 100$ and -1.09 at $t = 2 \times 10^4$, and more detailed fits show that the density is varying as $(t \ln t)^{-1}$. This is a consequence of the slow recovery of the superfluid density after the quench (bottom curve of Fig. 2b), which has a logarithmic approach to its equilibrium value. The substantial drop in the superfluid fraction just after the quench (from an initial value 0.89) is due to the increased polarizability of the pairs at low temperature. Their density is unchanged immediately after the quench, but

they screen more effectively due to the lack of thermal fluctuations.




FIG. 2: a) Decay of the pair density (in units a_o^{-2}) and b) recovery of the superfluid fraction (evaluated at $r = R$) in time after a quench to $0.1 T_{KT}$ from different starting temperatures T_i .

Quenches from starting temperatures below T_{KT} also display power-law variation of the vortex density, since the KT transition is a line of critical points. The plots in Fig. 2 show this for a series of initial temperatures between $0.8 T_{KT}$ and T_{KT} . At the lower starting temperatures the superfluid density recovers more quickly, and the vortex decay then varies accurately with the form $t^{-z/2}$, where the constant z agrees with the scaling prediction of Minnhagen and co-workers for the dynamic exponent [11],

$$z_{scale} = 4 \frac{\sigma_s(T)}{\sigma_s(T_{KT})} \frac{T_{KT}}{T} - 2 \quad , \quad (5)$$

Figure 3 shows the values of z extracted from the slopes of the curves in Fig. 2a, which are compared with z_{scale} , with a dynamic exponent predicted in Ref. [12], and with simulation results. It has recently been shown [8] that the simulations of the vortex dynamics can be misleading, since different results for z are obtained depending on the boundary conditions used: periodic boundary conditions (PBC) yield [7] only the spin-wave result $z = 2$ at all temperatures, while fluctuating-twist boundary conditions (FTBC) give z_{scale} [8]. Our results

in agreement with z_{scale} (except near T_{KT} where the logarithmic corrections enter) provide further support that this is the correct exponent characterizing the vortex dynamics. Unlike the simulations our formulation does not contain spin waves, and this allows a direct probe of the vortices, without their behavior being obscured by the slower dynamics of the spin waves.




FIG. 3: Dynamic exponent z calculated from the slope of the curves in Fig. 2a, compared with various scaling theories. The PBC simulations are from Ref. [7], the FTBC from the relaxational dynamics of Ref. [8].

Bray and co-workers [10] have predicted the appearance of a logarithmic term in the decaying vortex density for quenches starting above T_{KT} , but with a form $\ln t/t$ that differs from ours. In their calculation they assume, however, that the superfluid density is a constant in time. As seen in Fig. 2b this may be a reasonable approximation for quenches starting well below T_{KT} , but it appears to fail near T_{KT} because of the logarithmic recovery of the superfluid density. The log term in our vortex density arises entirely from this variation. In Ref. [10] the appearance of the logarithm is ascribed solely to the presence of free vortices, and this is cited as a violation of dynamic scaling. This is not the case in our calculation, where no free vortices at all are present. It appears that the development of the log term only depends on a sufficient density of pairs of large separation; for quenches starting from $T/T_{KT} = 0.98$ we find that the deviations from the power-law decay are already strong, but not quite the logarithmic behavior found right at T_{KT} .

Figure 4 illustrates the approach to equilibrium following an instantaneous quench from T_{KT} to $0.75 T_{KT}$. In the distribution function the smallest pairs quickly come to thermal equilibrium (the dashed line) at the new temperature, while the larger pairs only slowly readjust. The vortex density levels off near the equilibrium value, since it is dominated by the smaller pairs. The superfluid density, determined by the larger pairs, recovers more slowly, although because of the higher temperature its initial reduction is considerably smaller than that in Fig. 2b.




FIG. 4: a) Pair distribution function and b) vortex density and superfluid fraction (dotted curve) versus time for a quench from T_{KT} to $0.75 T_{KT}$.

From Fig. 2a it is clear that no vortices are being "created" in the instantaneous quenches from T_{KT} and below; there is only decay of the existing thermal vorticity. We have looked at the effect of the quench time on this decay, shown in Fig. 5. This shows the vortex density and superfluid fraction for the case of a linear ramp of the temperature from T_{KT} to $0.1 T_{KT}$ in a time τ_Q . We assume for convenience that D is independent of temperature, although in a real helium film it would vary during such a finite-time quench [16]. It can be seen in Fig. 5a that for a finite quench time the vortex density decreases more slowly than for the instantaneous quench (dashed curve), since the system spends more time at higher temperatures where the thermal density is higher. As the temperature falls and the smaller pairs that had been able to stay in quasi-equilibrium start to annihilate, the more sluggish larger pairs begin to dominate, and the curves revert to the instantaneous-quench result at long times. The slight wiggles in the curves are due to changes in the superfluid fraction, shown in Fig. 5b. The fraction begins to increase slightly as the density of the smaller quasi-equilibrium pairs decreases with the decreasing temperature, but finally the increasing polarizability of the larger pairs overcomes this, driving down the superfluid fraction. Since there are now more pairs remaining than in the instantaneous quench the fraction drops even below the value for that case, and then rejoins it at long times as the smaller pairs decay completely and the larger pairs dominate.

The behavior of the vortex density in Fig. 5 is the opposite




FIG. 5: Vortex density and superfluid fraction versus time for quenches from T_{KT} to $0.1 T_{KT}$ in time τ_Q (in units of the diffusion time).

of the Kibble-Zurek predictions of Ref. [1], which postulate that the faster the quench is performed the more vortices will be created. Since these theories do not take into account the thermal vortices it is unclear how any extra vortices would be formed. Of course, our calculations cannot be extended to starting temperatures above T_{KT} , since the KT recursion relations become invalid. We doubt, however, that the additional thermal vorticity generated at higher temperatures will do anything but add smoothly to the decaying curves of Fig. 5. The 2D quench simulations from above T_{KT} [6] show no evidence of any additional vortex creation. Even simulations of quenched 3D transitions show (in Fig. 1 of Ref. [17]) only a smooth and monotonic decay of the thermal vorticity starting from above T_c , with no extra vorticity being generated.

In summary, an analytic formulation of 2D quenched superfluid transitions shows a power-law decay of the vortex density with time, in agreement with the predictions for the dynamic scaling exponent of Minnhagen and co-workers. Quenches starting near T_{KT} show an additional logarithmic variation resulting from the slow recovery of the superfluid density to its equilibrium value. The strong depression of the superfluid density following a quench has not been previously been taken into account, though it plays a major role in the quench dynamics, as seen in Fig. 5. It will be important that any future simulations of quenched transitions (in both 2D and 3D) include computations of the superfluid fraction (the helicity modulus) as well as the vortex density, and also

study the effect of changing the boundary conditions. Experimental tests of these vortex dynamics should also be possible: measurements on Josephson-junction arrays have been able to determine $z = 1.98 \pm 0.03$ at T_{KT} [18], and experiments on liquid-crystal films are able to visualize the pair-defect annihilation process following a quench [19].

This work is supported by the National Science Foundation, DMR 97-31523.

REFERENCES

- [1] A. Gill and T. Kibble, J. Phys. A **29**, 4289 (1996); W. Zurek, Phys. Rep. **276**, 177 (1996). Extension of this theory to 2D superfluids is found in A. Yates and W. Zurek, Phys. Rev. Lett. **80**, 5477 (1998).
- [2] M. Dodd *et al.*, Phys. Rev. Lett. **81**, 3703 (1998).
- [3] Very different experiments in ^3He have claimed results consistent with the Kibble-Zurek predictions of Ref. 1: V. Ruutu *et al.*, Phys. Rev. Lett. **80**, 1465 (1998); C. Bäuerle *et al.*, Nature **382**, 332 (1996).
- [4] H. Chu and G. A. Williams, Physica B **284-288**, 115 (2000).
- [5] G. A. Williams, J. Low Temp. Phys. **93**, 1079 (1993); G. A. Williams, Phys. Rev. Lett. **82**, 1201 (1999).
- [6] R. Loft and T. DeGrand, Phys. Rev. B **35**, 8528 (1987); M. Mondello and N. Goldenfeld, Phys. Rev. A **42**, 5865 (1990); B. Yurke, A. Pargellis, T. Kovacs, and D. Huse, Phys. Rev. E **47**, 1525 (1993).
- [7] H. Luo and B. Zheng, Mod. Phys. Lett. **11**, 615 (1997).
- [8] B. Kim, P. Minnhagen, and P. Ollson, Phys. Rev. B **59**, 11506 (1999); L. Jensen, B. Kim, and P. Minnhagen, Phys. Rev. B **61**, 15412 (2000).
- [9] A. Rutenberg and A. Bray, Phys. Rev. E **51**, R1641 (1995).
- [10] A. Bray, A. Briant, and D. Jervis, Phys. Rev. Lett. **84**, 1503 (2000).
- [11] P. Minnhagen, O. Westman, A. Jonsson, and P. Olsson, Phys. Rev. Lett. **74**, 3672 (1995).
- [12] V. Ambegaokar, B. Halperin, D. Nelson, and E. Siggia, Phys. Rev. B **21** 1806 (1980).
- [13] J.M. Kosterlitz and D. Thouless, J. Phys. C **6**, 1181 (1973); J.M. Kosterlitz, J. Phys. C **7**, 1046 (1974).
- [14] D. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).
- [15] In this formulation it is not necessary to follow the complex motion and reconnection of the pairs with each other, as such motion is reflected statistically in the change of the distribution function after each time step. Our recombination process at a_o is the same as the “bulk recombination” process discussed by other authors who follow the collision of individual pairs [12].
- [16] H. Yano, T. Jocha, and N. Wada, Phys. Rev. B **60** 543 (1999).
- [17] N. Antunes, L. Bettencourt, and W. Zurek, Phys. Rev. Lett. **82**, 2824 (1999).
- [18] T. Shaw *et al.*, Phys. Rev. Lett. **76**, 2551 (1995).

- [19] C. Muzny and N. Clark, Phys. Rev. Lett. **68**, 804 (1992);
N. Clark, private communication.